arXiv:1211.5061v1 [hep-ph] 21 Nov 2012

Probing small $x(g)$ region with the LHeC based γp colliders

U. Kaya[1], S. Sultansoy[1,2], G. Unel[3]
[1]TOBB Univesity of Economics and Technology, Ankara, Turkey
[2]ANAS Institute of Physics, Baku, Azerbaijan
[3]UC Irvine,USA

Abstract

Potential of the LHeC for probing small $x(g)$ region via $c\bar{c}$ and $b\bar{b}$ production have been investigated. Obtained results clearly show the advantage of γp collider option. Measurement of $x(g)$ down to 3×10^{-6} seems to be reachable which is two order smaller than HERA coverage.

1 Introduction

The problem of precise measurement of parton distribution functions (PDF) is yet to be solved for the energy scales relevant for LHC results. On the other hand, precision knowledge on parton distribution of small x_B and sufficiently large Q^2 is crucial for enlightening of QCD basics at all levels from partons to nuclei. One of the measurements needed is the gluon PDF for low momentum fraction: small $x(g)$. The last machine which has probed $x(g)$ was HERA which had a reach of about $x(g) > 10^{-4}$. Large Hadron-Electron Collider (LHeC) project [1] - the most powerful microscope ever designed - will provide a uniqe opportunity to probe extremely small $x(g)$ region.

Within the linac-ring option of the LHeC the proton beam from LHC can be hit with a high energy electron or photon beam. This option gaining wider acceptance among the high energy physics community [2] . The photons may be virtual ones from the electron beam resulting in a typical DIS event or these can be real photons originating from the Compton Back Scattering process. In the latter case, the photon spectrum consists of the high energy photons peaking at about 80% of the electron beam energy on the continum of Weizsacker-Williams photons. The following study aims to investigate the fesibility of a small $x(g)$ measurement with such a machine. Main parameters of ep and γp options of the LHeC are presented in section 2. Section 3 is devoted to investigation of small $x(g)$ region using the processes $\gamma p \to c\bar{c}X$ and $\gamma p \to b\bar{b}X$. The generator level results are obtained using CompHEP [3] software package. Comparision with processes $ep \to ec\bar{c}X$ and $ep \to eb\bar{b}X$ shows an obvious advantage of the LHeC γp option. Finally, in section 4 summary of obtained results are presented together with some recommendations.

2 Main parameters of ep and γp options of the LHeC

It should be emphasized that real γp collisions can be achieved only on the base of linac ring type ep colliders (see review [4] for history and status of linac-ring type collider proposals). Real γ beam for γp collider [5] will be produced using the Compton back scattering of laser beam off the high energy electron beam [6] . Possible application of this mechanism to ring-ring type ep colliders result in negligible γp luminosities $L_{\gamma p} < 10^{-7} L_{ep}$.

Currently, two versions for the ep option of the LHeC are under consideration: multi-passed energy recovery linac with $L_{ep} = 10^{33} cm^{-2}s^{-1}$ and pulsed single pass linac with $L_{ep} = 10^{32} cm^{-2}s^{-1}$. In the first case $E_e = 60\,GeV$ has been choosen, higher energies are not available because of the synchrotron radiation in arcs. In the second case beam energies above $140\,GeV$ will be available [1]. Main parameters of ep collisions are presented Table 1.

Table 1: Main parameters of ep collisions.

	E_e, GeV	E_p, TeV	$\sqrt{s}$, TeV	L, $cm^{-2}s^{-1}$
ERL	60	7	1.30	10^{33}
Pulsed 1	60	7	1.30	9×10^{31}
Pulsed 2	140	7	1.98	4×10^{31}

Concerning γp options luminosity of γp collisions will be close to luminosity of ep collisions for pulsed single straight linac. In the ERL case, $L_{\gamma p}$ will be 10 times lower than L_{ep} because energy recovering does not work after Compton back scattering.

3 Pair production of $c\bar{c}$ and $b\bar{b}$ at the LHeC

The final states that can be easily distinguished from the background events and that would give a good measure of the $x(g)$ are $eg \to eq\bar{q}$ and/or $\gamma g \to q\bar{q}$ where the gluon (g) is from the LHC protons, electrons and photons are from a new accelerator (namely, an electron linac providing beams tangential to the LHC) to be build and the q stands for a heavy quark flavour, such as b quark and possibly c as well. The b quark final states are easier to idenify due to b-tagging possibility using currently available technologies: for example, ATLAS silicon detectors have 70% b-tagging efficiency. In the Table 2 we present cross sections for heavy quark pair production via DIS, quasi real photons (WWA) and Compton Back Scattering (CBS) photons at the LHeC with $E_e = 60\,GeV$ and $E_e = 140\,GeV$. For comparison, we also give values for DIS and WWA processes at HERA. It is seen that WWA quasi real photons are adventageous comparing to DIS and CBS photons are advantageous comparing to WWA. All numerical calculations are performed using CompHep [3] with CTEQ6L1 [7] PDF distrubutions. In Figure 1, differantial cross section depending on the $x(g)$ have been shown for WWA photons at HERA and the LHeC. As expected, LHeC will give opportunity to investigate an order smaller $x(g)$ than HERA.

Table 2: Heavy quark pair production cross sections for via DIS, WWA, and CBS.

	$b\bar{b}\,(pb)$			$c\bar{c}\,(pb)$		
Machine	DIS	WWA	CBS	DIS	WWA	CBS
HERA	6.07×10^2	4.57×10^3	-	4.66×10^4	7.29×10^5	-
LHeC 1($E_e = 60\,GeV$)	4.26×10^3	2.99×10^4	2.41×10^5	2.38×10^5	3.44×10^6	2.38×10^7
LHeC 2($E_e = 140\,GeV$)	7.07×10^3	4.91×10^4	3.70×10^5	3.72×10^5	5.27×10^6	3.46×10^7

The advantage of the CBS photons becomes even more obvious if one analyze $x(g)$ distribution of differantial cross sections for CBS, WWA and DIS. In Figure 2, we show the $d\sigma/dx(g)$ at the LHeC 1 for $c\bar{c}$ production. It is seen that CBS at small $x(g)$ region provides more than one (two) order higher cross sections comparing to WWA (DIS). For example, differantial cross section of $c\bar{c}$ pair production at the LHeC 1 achieves maximum value $94\,\mu b$ at $x(g) = 1.44 \times 10^{-5}$ for CBS, whereas maximum value for WWA and DIS are $4\,\mu b$ at $x(g) = 1.54 \times 10^{-5}$ and $0.15\,\mu b$ at $x(g) = 3.89 \times 10^{-5}$, respectively. Similar distrubutions for $b\bar{b}$ at LHeC 1, $c\bar{c}$ at LHeC 2 and $b\bar{b}$ at LHeC 2 are shown in Figures 3, 4 and 5, respectively. Maximum values of differantial cross sections and corresponding $x(g)$ values for DIS, WWA, and CBS at the LHeC 1 (2) are given in the Table 3 (4). The advantage of CBS due to large cross section is obvious.

Table 3: Maximum values of differantial cross sections and corresponding $x(g)$ values for DIS, WWA, and CBS at the LHeC 1.

	$c\bar{c}$		$b\bar{b}$	
	$d\sigma/dx$	x	$d\sigma/dx$	x
DIS	$0.15\,\mu b$	3.89×10^{-5}	$0.47\,nb$	1.99×10^{-4}
WWA	$4.0\,\mu b$	1.54×10^{-5}	$5.02\,nb$	1.25×10^{-4}
CBS	$94\,\mu b$	1.44×10^{-5}	$117\,nb$	1.23×10^{-4}

Table 4: Maximum values of differantial cross sections and corresponding $x(g)$ values for DIS, WWA, and CBS at the LHeC 2.

	$c\bar{c}$		$b\bar{b}$	
	$d\sigma/dx$	x	$d\sigma/dx$	x
DIS	$0.44\,\mu b$	1.54×10^{-5}	$1.73\,nb$	9.12×10^{-5}
WWA	$13.2\,\mu b$	6.45×10^{-6}	$17\,nb$	5.88×10^{-5}
CBS	$312\,\mu b$	6.02×10^{-6}	$408\,nb$	5.01×10^{-5}



Figure 1: The x(g) reach and differential cross sections in $c\bar{c}$ (left) and $b\bar{b}$ (right) final states for the HERAand the LHeC.



Figure 2: Differential cross sections for $c\bar{c}$ final states produced via CBS, WWA and DIS at the LHeC 1.



Figure 3: Differential cross sections for $b\bar{b}$ final states produced via CBS, WWA and DIS at the LHeC 1.



Figure 4: Differential cross sections for $c\bar{c}$ final states produced via CBS, WWA and DIS at the LHeC 2.



Figure 5: Differential cross sections for $b\bar{b}$ final states produced via CBS, WWA and DIS at the LHeC 2 .

The angular dependency of the relevant processes is important to estimate the necessary η coverage of the detector to be built and also to estimate the eventual electron machine selection. For illustration we consider $d\sigma/d\theta$ distrubution where θ is the angle between c (b) quark and proton beam direction. These distributions for CBS at the LHeC 1 and LHeC 2 are presented in Figures 6 and 7, respectively. In Table 5 we present reachable $x(g)$ for diffrent θ coverage. One can notice that even for an angular loss of about 5 degrees, there is considerable drop in both the cross section and in the $x(g)$ reach. This effect can be understood by considering the η depence of the heavy quark pair production cross section in γp collisions which are shown in Figure 8 and 9. The vertical solid line is representative for a 1 degree, the dashed line for a 5 degree and the dot-dashed line is for 10 degree detector. Therefore, in order to have a good experimental reach the tracking should have an η coverage up to 5.

Table 5: Reachable $x(g)$ for different θ coverage.

	LHeC 1		LHeC 2	
θ	$c\bar{c}$	$b\bar{b}$	$c\bar{c}$	$b\bar{b}$
0-180	7.94×10^{-6}	6.91×10^{-5}	3.16×10^{-6}	3.02×10^{-5}
1-179	8.31×10^{-6}	6.91×10^{-5}	3.36×10^{-6}	4.36×10^{-5}
5-175	1.44×10^{-5}	7.94×10^{-5}	1.20×10^{-5}	4.78×10^{-5}
10-170	2.39×10^{-5}	1.0×10^{-4}	2.28×10^{-5}	7.58×10^{-5}



Figure 6: The effect of angular reach for $c\bar{c}$ (left) and $b\bar{b}$ (right) final states produced via CBS at the LHeC 1.



Figure 7: The effect of angular reach for $c\bar{c}$ (left) and $b\bar{b}$ (right) final states produced via CBS at the LHeC 2.



Figure 8: The η dependency of the $c\bar{c}$ (left) and $b\bar{b}$ (right) production cross section via CBS at the LHeC 1.



Figure 9: The η dependency of the $c\bar{c}$ (left) and $b\bar{b}$ (right) production cross section via CBS at the LHeC 2.

4 Conclusions

Measurements of $x(g)$ down to 3×10^{-6} seems to be reachable in γp collisions which is two order smaller than HERA coverage. These collisions provide higher cross section and better $x(g)$ reach with respect to the ep collisions with the same electron beam energy. For the low $x(g)$ region, the enhancement factor compared to the DIS ep collisions is about 700 for $c\bar{c}$ final states and about 230 for $b\bar{b}$ final states at the LHeC 2. Therefore, for the final states with heavy quarks even if the γp luminosity is 10 times less than ep luminosity, the expected number of events would be 70 and 20 times higher, respectively. The enhancement factor compared to WWA ep collisions is about 24 for both final states. The angular sensitivity is very important for smallest $x(g)$ reach for either e or γ beams, coverage up to $\eta = 5$ is required. This coverage is already achieved at the ATLAS and CMS with using forward detector components.

References

[1] J.L.Abelleira Fernandez , C.Adolphsen , A.N.Akay et al., J. Phys. G: Nucl. Part. Phys. 39 (2012) 075001.

[2] J.L.Abelleira Fernandez , C.Adolphsen , P.Adzic et al., arXiv:1211.4831v1 [hep-ex] 20 Nov 2012.

[3] E. Boos et al. (CompHEP Collaboration), Nucl. Instrum. Meth. A 534, 250 (2004)

[4] A. N. Akay, H. Karadeniz, and S.Sultansoy, Int. J. Mod. Phys. A 25, 4589 (2010).

[5] S. I. Alekhin et al., Int. J. Mod. Phys. A 6, 21 (1991); A. K. Ciftci et al., Nucl. Instrum. Methods Phys. Res., Sect. A 365, 317 (1995); A. K. Ciftci, S. Sultansoy, and O. Yavas, Nucl. Instrum. Methods Phys. Res.,Sect. A 472, 72 (2001); H. Aksakal et al., Nucl. Instrum. Methods Phys. Res., Sect. A 576, 287 (2007).

[6] I. F. Ginzburg et al., Nucl. Instrum. Methods Phys. Res., Sect. A 219, 5 (1984); V. I. Telnov, Nucl. Instrum. Methods Phys. Res., Sect. A 355, 3 (1995).

[7] J. Pumplin et al., JHEP 0207 (2002) 012.